



Search for $\Lambda_b^0 \rightarrow pK$ and $p\pi$ decays

The CDF Collaboration
URL <http://www-cdf.fnal.gov>
(Dated: July 7, 2005)

This note describes a search for $\Lambda_b^0 \rightarrow pK$ and $\Lambda_b^0 \rightarrow p\pi$. The analysis is based on an integrated luminosity of 193 pb^{-1} collected with the CDF II detector. The preliminary result is an upper limit on the Branching Ratio. We set the upper limits on the branching fraction $\mathcal{B}(\Lambda_b^0 \rightarrow ph)$, where h is K or π , of 2.3×10^{-5} at 90% C.L. and 2.9×10^{-5} at 95% C.L.

Preliminary Results for Summer 2004 Conferences

I. INTRODUCTION

Detection of Λ_b^0 charmless decays into hadrons is one of the challenging goals of CDF II [1]. In those decays, a quite large direct CP violation is expected [2].

This note describes a search for the decays $\Lambda_b^0 \rightarrow pK$ and $\Lambda_b^0 \rightarrow p\pi$ in $\bar{p}p$ collisions at $\sqrt{s} = 1.96$ TeV with the CDF detector at the Fermilab Tevatron. Such decays have not been seen up to now. The current upper limits on the Branching Ratios are $50. \times 10^{-6}$ on each of the decay modes [3]. Theoretical predictions are between 10^{-6} and $2. \times 10^{-6}$ [4].

The Branching Ratios $BR(\Lambda_b^0 \rightarrow hh)$ are normalized to the Branching Ratio $BR(B_d^0 \rightarrow K\pi) = (1.85 \pm 0.11) \times 10^{-5}$ [5].

The relationship between the Branching Ratios and the number of observed events is the following:

$$\frac{\epsilon_\Lambda f_\Lambda BR(\Lambda_b^0 \rightarrow hh)}{\epsilon_B f_B BR(B_d \rightarrow K\pi)} = \frac{N(\Lambda_b^0 \rightarrow hh)}{N(B_d \rightarrow K\pi)}$$

where:

- N is the number of observed events (Λ_b^0 or B_d);
- ϵ_Λ (ϵ_B) is the total efficiency (trigger, reconstruction and analysis) for observing a Λ_b^0 (B^0); The efficiency ϵ_Λ is assumed to be the same for the two decay modes pK and $p\pi$;
- f_Λ (f_d) is fraction of b -quarks that hadronize into Λ_b^0 (B_d^0). We use the following values [5]: $f_\Lambda = 0.094 \pm 0.017$, $f_B = 0.397 \pm 0.010$, $f_\Lambda/f_B = 0.25 \pm 0.04$

The number of observed $B_d \rightarrow K\pi$ decays is a fraction of the total number of $B \rightarrow h^+h^-$:

$$f = N(B_d \rightarrow K\pi)/N(B \rightarrow h^+h^-)$$

Therefore

$$BR(\Lambda_b^0 \rightarrow hh) = \frac{N(\Lambda_b^0 \rightarrow hh)}{A}$$

where

$$A = \frac{\epsilon_\Lambda f_\Lambda f N(B \rightarrow h^+h^-)}{\epsilon_B f_B BR(B_d \rightarrow K\pi)}$$

We call A the acceptance.

II. DATA SAMPLE & EVENT SELECTION

The analysis is performed on data taken with the CDFII detector between February 2002 and September 2003, corresponding to an integrated luminosity of 193 pb^{-1} .

The data are collected with the “Two-Track Trigger”, which selects events containing track pairs coming from secondary vertexes.

CDF uses a three-level trigger. At first level, the XFT (eXtremely Fast Tracker) reconstructs and selects tracks with transverse momentum $p_T > 2 \text{ GeV}/c$.

At the second level the SVT (Silicon Vertex Tracker) measures the track impact parameters (d_0) and selects tracks pairs coming from secondary vertexes by cutting on d_0 .

At the third level a complete event reconstruction is performed, and the level 1 and level 2 requirements are confirmed.

In order to look for $\Lambda_b^0 \rightarrow pK$ and $p\pi$ decays, the invariant mass of track pairs is calculated by assigning the mass of the π to both tracks. The reason is that such assignment maximizes the separation between the peaks of the $B \rightarrow hh$ and the Λ_b^0 decays in the mass spectrum.

A. Optimized Search

The selection optimization has been performed using the estimator defined in [6]. This estimator provides optimal cuts which are good for signal measurements and for setting an upper limit on it.

The optimization has been performed hiding the signal region in the invariant mass spectrum and using only events with even trigger numbers, i.e. 50% of the total sample. In fact, we want to avoid using the same sample for both optimization and background calculation, since we can introduce a statistical bias due to fluctuations.

For each set of cuts, the value of the background has been obtained by fitting the sidebands in the invariant mass spectrum and integrating the fitted function in the signal region.

The reconstruction efficiency for the signal has been evaluated using a Monte Carlo.

III. BACKGROUNDS

The background in this analysis is given by track pairs which accidentally have an invariant mass close to the mass of the Λ_b^0 (“combinatorial background”) and the tail from the $B \rightarrow hh$ signal.

The background has been calculated by fitting the invariant mass spectrum and interpolating in the signal region. Only events with odd trigger numbers were used.

In order to evaluate the systematic uncertainty due to the shape of the background, different functions have been used. Such a systematic affects the number of $B \rightarrow hh$ events used for normalization.

The value for the expected background in the signal region is:

$$b = 772 \pm 31$$

where the error includes both the statistical and systematic errors.

IV. SYSTEMATIC UNCERTAINTIES

Systematic uncertainties on the acceptance come from the following sources:

- shape of the background in the signal region of the invariant mass spectrum (it affects the expected background);
- the number of $B \rightarrow hh$ reconstructed decays is also affected by the shape of the background;
- relative Branching Fractions of $\Lambda_b^0 \rightarrow pk$ and $\Lambda_b^0 \rightarrow p\pi$, since the two decay modes have a different reconstruction efficiency;
- difference between the invariant masses predicted by Monte Carlo and reconstructed in the data;
- difference between the invariant mass resolutions predicted by Monte Carlo and reconstructed in the data;
- uncertainties in the B and Λ_b^0 lifetimes;
- different Level 1 trigger efficiencies for Protons, Kaons and Pions, because of the different specific ionization;
- different p_T spectra for B and Λ_b^0 s;
- uncertainty in the Branching Ratio $BR(B \rightarrow K\pi)$;
- uncertainty in the production fractions of B and Λ_b^0 .

In table I all the systematic errors are listed.

Affected qty.	Source	Syst. Error
$B \rightarrow h^+ h'^- \text{ yield}$	Bkg. shape	5.7%
Bkg. estimate	Bkg. shape	3.3%
$\epsilon_{\Lambda_b^0} / \epsilon_B$	$(\Lambda_b^0 \rightarrow p\pi) / (\Lambda_b^0 \rightarrow pK)$	3.5%
	Window position	1.2%
	Window width	9%
	Lifetime	3.6%
	proton's trigger efficiency	6%
	$p_T(\Lambda_b^0)$	17%
	Overall	21%
$\mathcal{B}(B_d \rightarrow K\pi)$		5.9%
f_Λ / f_d		16%

TABLE I: Summary of the systematics.

$N(B \rightarrow hh)$	726 ± 82
$\epsilon_B / \epsilon_\Lambda$	1.77 ± 0.37

TABLE II: Measured quantities that enter in the Λ_b^0 acceptance.

V. RESULTS

In figure 1 we show the full mass spectrum with the search window. The total number of events in the signal region of the mass spectrum is

$$N = 767$$




FIG. 1: Mass distribution obtained including all the events. The dashed curve shows the fitted function in the part of the mass range that was excluded from the fit. The scales of the Monte Carlo distributions of the two signal decay modes are arbitrary. The peak in the data is given by the $B \rightarrow h^+ h^-$ events.

The quantity

$$f = \frac{N(B \rightarrow K\pi)}{N(B \rightarrow h^+h^-)} = 0.59 \pm 0.04$$

has been calculated using the fitting procedure used in [7].

The ratio between the number of $\Lambda_b^0 \rightarrow hh$ and the branching ratio $BR(\Lambda_b^0 \rightarrow hh)$ depends on the number of events of $B \rightarrow hh$ and the relative efficiency $\epsilon_\Lambda/\epsilon_B$. These values are reported in table II. The measured value of the ratio is:

$$A = \frac{N(\Lambda_b^0 \rightarrow hh)}{BR(\Lambda_b^0 \rightarrow hh)} = (3.2 \pm 1.0) \cdot 10^6$$

The error includes both statistical and systematic errors.

Using a Bayesian method uniform prior distribution, we find that the upper limit on the number of signal events s and on $BR(\Lambda_b^0 \rightarrow ph)$ are respectively 75 and 2.3×10^{-5} at 90% C.L and 97 and 2.9×10^{-5} at 95% C.L.

Acknowledgments

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Particle Physics and Astronomy Research Council and the Royal Society, UK; the Russian Foundation for Basic Research; the Comisión Interministerial de Ciencia y Tecnología, Spain; in part by the European Community's Human Potential Programme under contract HPRN-CT-2002-00292; and the Academy of Finland.

-
- [1] F. Abe, et al., Nucl. Instrum. Methods Phys. Res. A **271**, 387 (1988); D. Amidei, et al., Nucl. Instrum. Methods Phys. Res. A **350**, 73 (1994); F. Abe, et al., Phys. Rev. D **52**, 4784 (1995); P. Azzi, et al., Nucl. Instrum. Methods Phys. Res. A **360**, 137 (1995); The CDFII Detector Technical Design Report, Fermilab-Pub-96/390-E
 - [2] I. Duniety, Z. Phys. C **56** (1992) 129.
 - [3] D. Buskukic et al., Phys. Lett. **B 384** (1996) 471-480.
 - [4] R. Mohanta et al., Phys. Rev. **D 63**:074001 (2001).
 - [5] *Review of Particle Physics*, S. Eidelman *et al.*, Phys. Lett. **B592** (2004).
 - [6] G. Punzi, Optimization of searches for new signals, physics/0308063, <http://www.arxiv.org/ps/physics/0308063>.
 - [7] G. Punzi, in *Proceedings 32nd International Conference on High-Energy Physics (ICHEP 04)*, (2004)